



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

C being a constant, individual for each liquid; being taken at 8.7 for water and m at 18 the last equation yields

$$T_x = \frac{m}{1.49} (p_x v_x - 8.7) = 12.1 (p_x v_x - 8.7).$$

Since we calculate from tables of properties of saturated vapor of water that *pv* at 273° absolute amounts to 31 calories per kilogram the above equation for T_x gives

$$12.1 (31-8.7) = 270^\circ \text{ absolute}$$

at 313 degrees at which *pv* is 34.6 cal. we find

$$12.1 (34.6-8.7) = 313.3^\circ \text{ absolute.}$$

At 473 degrees absolute *pv* is equal 47.9 calories and

$$12.1 (47.9-8.7) = 473.5^\circ \text{ absolute.}$$

Similar agreement is found for other vapors by inserting the correct value for constant as long as no polymerisation in the liquid takes place.

(1) *Molecular state of water vapor*; (2) *Vapor pressure depression equation for dilute aqueous solutions*: JAMES KENDALL.

Size and behavior of suspended smoke particles: R. E. WILSON.

Influence exerted by antagonistic electrolytes on the electrical resistance and permeability of emulsion membranes: G. H. A. CLOWES.

The exact determination of molecular weights by the boiling point method: E. M. WASHBURN.

Solubility of strontium nitrate in anhydrous alcohol in alcohol containing small per cent. of water: C. W. FOULK.

(1) *Influence of the age of ferric arsenate on its peptization*; (2) *Syneresis of silicic acids gels*: H. N. HOLMES.

A study of the lowering of vapor pressure of water produced by absorbed KCl: B. F. LOVELACE, J. C. W. FRAZER, V. B. SEASE.

A study of the lowering of vapor pressure of water produced by absorbed mannite: J. C. W. FRAZER, B. F. LOVELACE, T. H. ROGERS.

The volume and surface of the pores in charcoal and the compression of adsorbed substances: W. D. HARKINS and D. T. EWING.

An electromagnetic and valence hypothesis of heterogeneous equilibrium in adsorption: W. D. HARKINS.

DIVISION OF WATER, SEWERAGE AND SANITATION

Robert Spurr Weston, *Chairman*

W. W. Skinner, *Secretary*

Determination of bromid in mineral waters and brines: W. W. SKINNER and W. F. BAUGHMAN.

Colorimetric methods for the determination of bromin give satisfactory results only when small quantities of bromin are to be determined. The method proposed for the determination of bromids in the presence of chlorids is the oxidation of the bromids and removal of the liberated bromid by steam distillation or by aspiration. The method depends upon the use of chromic acid for oxidation of the bromid. Chromic acid in concentrated solution liberates bromin from bromids quantitatively at room temperature and the bromin may be removed by aspiration. It liberates only a trace of chlorin from chlorids, forming probably chromic chlorid which remains in solution. When chromic acid acts on a solution of chlorids and bromids, some chlor-bromid is formed which is removed with the bromin by aspiration. The liberated bromin and the chlorin in the first aspiration is collected in a solution of sodium sulphite and sodium carbonate, which is evaporated to dryness and again submitted to the treatment with chromic acid and aspirated the second time. The double aspiration gives very accurate results.

Certain war gases and health: CHARLES BASKERVILLE. Evidence has been collected from all the chlorine producing plants and many works and arsenals where chlorine was used. Preponderating evidence favors the conclusion that chlorine exerts a preventative influence against influenza. The evidence is not conclusive, however, as contrary data were obtained from some plants. The contradictions may possibly be harmonized on the basis of concentration, the more dilute up to limits the more effective. Small amounts of bromine in the air appear to prevent influenza completely.

CHARLES L. PARSONS,
Secretary

SCIENCE

A Weekly Journal devoted to the Advancement of Science, publishing the official notices and proceedings of the American Association for the Advancement of Science

Published every Friday by

THE SCIENCE PRESS

LANCASTER, PA.

GARRISON, N. Y.

NEW YORK, N. Y.

Entered in the post-office at Lancaster, Pa., as second class matter